Neuroscience Letters, 165 (1994) 227–229
Elsevier Science Ireland Ltd.

AUTHOR INDEX

Achaaban, M.R., see Wells, T., 93
Alhayek, G., see Dehaut, F., 59
Amaducci, L., see Mortilla, M., 45
Anadón, R., see Yáñez, J., 63
Anderson, J.H., see Kaufman, G.D., 144
Aparicio, J.J., see Blanco, P., 5
Arilla, E., see López-Sañudo, S., 27
Arndt, J.O., see Holthusen, H., 71
Atlas, D., see Avidor, T., 133
Avidor, T., Clementi, E., Schwartz, L. and Atlas, D., Caffeine-induced transmitter release is mediated via ryanodine-sensitive channel, 133
Avila, J., see Szücs, K., 175

Baranauskas, G., see Fisher, N.D., 84
Baron, R. and Saguer, M., Axon-reflex reactions in affected and homologous contralateral skin after unilateral peripheral injury of thoracic segmental nerves in humans, 97
Behnisch, T. and Reymann, K.G., 2,3-Diphosphoglyceric acid blocks long-term potentiation of excitatory postsynaptic currents in hippocampal CA1 neurons of the rat, 23
Beitz, A.J., see Kaufman, G.D., 144
Berrino, L., see Maione, S., 164
Birch, P.J., see Smith, G.D., 79
Bitting, L., Sutin, E.L., Watson, F.L., Leard, L.E., O'Hara, B.F., Heller, H.C. and Kilduff, T.S., C-*fos* mRNA increases in the ground squirrel suprachiasmatic nucleus during arousal from hibernation, 117
Blanco, P., Huidobro, A., Villalba, M., Gómez-Puertas, P., Bogónez, E., Pereira, R., Martínez-Serrano, A., Aparicio, J.J. and Satrústegui, J., The activity of synaptosomal calcium channels is inversely correlated with working memory performance in memory impaired, aged rats, 5
Bogónez, E., see Blanco, P., 5
Bongrani, S., see Gandolfi, O., 113
Brady, L.S., see Young, J.K., 13
Brown, G.M., see Pang, C.S., 55
Bruni, A., see Mortilla, M., 45
Brysch, W., see Herms, J., 171
Budhram-Mahadeo, V., Lillycrop, K.A. and Latchman, D.S., Cell cycle arrest and morphological differentiation can occur in the absence of apoptosis in a neuronal cell line, 18
Bugnon, C., see Griffond, B., 89
Bunney, B.S., see Youngren, K.D., 41

Catt, K.J., see Krsmanovic, L.Z., 33
Chan, J.Y.H., see Tsai, H.-F., 137
Chan, S.H.H., see Fu, M.-J., 199
Chan, S.H.H., see Tsai, H.-F., 137
Chia, L.G., see Lee, E.H.Y., 149
Chino, Y.M., see Garraghty, P.E., 223
Chung, J.M., see Park, M.J., 125
Chung, K., see Park, M.J., 125
Clementi, E., see Avidor, T., 133
Colard, C., see Griffond, B., 89
Couratier, P., see Diop, A.G., 187

Dall'Olio, R., see Gandolfi, O., 113
De Cataldo, S., see Mortilla, M., 45
Dehaut, F., Haddad, K., Alhayek, G. and Pouplard-Barthelaix, A., Autoantibodies against H- and M-subunits of neurofilaments are induced by PC12 cell grafts or lesions into different sites of rat brain, 59
Deray, A., see Griffond, B., 89
Derouiche, A. and Ohm, T.G., Glutamine synthetase immunoreactivity in the human hippocampus is lamina-specific, 179
Diop, A.G., Lesort, M., Esclaire, F., Sindou, P., Couratier, P. and Hugon, J., Tetrodotoxin blocks HIV coat protein (gp120) toxicity in primary neuronal cultures, 187
Dohanics, J., Hoffman, G.E. and Verbalis, J.G., Hyponatremia inhibits sprouting of oxytocinergic axons following bilateral lesions of the paraventricular nucleus in rats, 51
Dombrádi, V., see Szücs, K., 175

Elliott, P.J., see Smith, G.D., 79
Esclaire, F., see Diop, A.G., 187

Farkas, R.H., Nakajima, S. and Nakajima, Y., Cultured neurons infected with an HSV-1-derived vector remain electrically excitable and responsive to neurotransmitter, 153
Fellmann, D., see Griffond, B., 89
Fisher, N.D., Baranauskas, G. and Nistri, A., Multiple types of tachykinin receptor mediate a slow excitation of rat spinal motoneurones in vitro, 84
Forsling, M.L., see Wells, T., 93
Fox, J.A., Novel ω-conopeptides reduce field potential amplitudes in the rat hippocampal slice, 157
Friedrich, P., see Szücs, K., 175
Fu, M.-J., Lin, T.C.Y., Lui, P.-W., Lee, T.-Y. and Chan, S.H.H., Involvement of $G_{o\alpha}$ subtype of guanine nucleotide-binding regulatory protein at the locus coeruleus in fentanyl-induced muscular rigidity in the rat, 199
Fujie, S., see Watanabe, K., 9
Fujii, T., Sakai, M. and Nagatsu, I., Immunohistochemical demonstration of expression of tyrosine hydroxylase in cerebellar Purkinje cells of the human and mouse, 161
Fujita, K., see Mogi, M., 208

Gandolfi, O., Dall'Olio, R., Rimondini, R., Bongrani, S. and Valsecchi, B., Repeated administrations of (−)-deprenyl increase [^{3}H]MK801 binding in rat brain and antagonize the impairment of passive avoidance conditioning induced by *N*-methyl-D-aspartate receptor antagonists, 113
Garraghty, P.E., Roe, A.W., Chino, Y.M. and Sur, M., Abnormal development of retinogeniculate X axons in strabismic cats: a possible substrate for visual dysfunction, 223
Gergely, P., see Szücs, K., 175
Gómez-Puertas, P., see Blanco, P., 5
Goso, C., see Szallasi, A., 101
Grecu, L., see Mortilla, M., 45
Greenberg, D.A., see von B. Ahern, K., 211
Griffond, B., Colard, C., Deray, A., Fellmann, D. and Bugnon, C., Evidence for the expression of dynorphin gene in the prolactin-immunoreactive neurons of the rat lateral hypothalamus, 89

Haddad, K., see Dehaut, F., 59
Harada, M., see Mogi, M., 208

Harrison, S.M., see Smith, G.D., 79
Hasegawa, T., see Yamada, K., 191
Headley, P.M., see Herrero, J.F., 37
Heller, H.C., see Bitting, L., 117
Herkenham, M., see Young, J.K., 13
Herms, J., Zurmöhle, U., Schlingensiepen, R., Brysch, W. and Schlingensiepen, K.-H., Developmental expression of the transcription factor *zif268* in rat brain, 171
Herrero, J.F., Headley, P.M. and Parsons, C.G., Memantine selectively depresses NMDA receptor-mediated responses of rat spinal neurones in vivo, 37
Hitosugi, H., see Kamei, J., 141
Hoffman, G.E., see Dohanics, J., 51
Holthusen, H. and Arndt, J.O., Nitric oxide evokes pain in humans on intracutaneous injection, 71
Hugon, J., see Diop, A.G., 187
Huidobro, A., see Blanco, P., 5

Inoue, Y., see Watanabe, M., 183
Iwamoto, Y., see Kamei, J., 141

Kamei, J., Iwamoto, Y., Hitosugi, H., Misawa, M., Nagase, H. and Kasuya, Y., Streptozotocin-induced diabetes selectively reduces antinociception mediated by μ_1-opioid receptors, but not that mediated by μ_2-opioid receptors, 141
Kasuya, Y., see Kamei, J., 141
Kaufman, G.D., Anderson, J.H. and Beitz, A.J., Hemilabyrinthectomy causes both an increase and a decrease in corticotropin releasing factor mRNA in rat inferior olive, 144
Kehl, S.J., Voltage-clamp analysis of the voltage-gated sodium current of the rat pituitary melanotroph, 67
Kilduff, T.S., see Bitting, L., 117
Koch, B., see Lutz-Bucher, B., 48
Koh, D.-S., Reid, G. and Vogel, W., Effect of the flavoid phloretin on Ca^{2+}-activated K^+ channels in myelinated nerve fibres of *Xenopus laevis*, 167
Krsmanovic, L.Z., Virmani, M.A., Stojilkovic, S.S. and Catt, K.J., Stimulation of gonadotropin-releasing hormone secretion by acetyl-L-carnitine in hypothalamic neurons and GT1 neuronal cells, 33
Kuo, J.S., see Lee, E.H.Y., 149
Kuo, T.B.J., see Tsai, H.-F., 137

Latchman, D.S., see Budhram-Mahadeo, V., 18
Leard, L.E., see Bitting, L., 117
Ledesma, M.D., see Szücs, K., 175
Lee, E.H.Y., Yang, Y.L., Chia, L.G. and Kuo, J.S., Interactions among deprenyl, clorgyline and 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine on catecholamines in mice, 149
Lee, T.-Y., see Fu, M.-J., 199
Lesort, M., see Diop, A.G., 187
Leyva, J., see Maione, S., 164
Lillycrop, K.A., see Budhram-Mahadeo, V., 18
Lin, T.C.Y., see Fu, M.-J., 199
López-Sañudo, S. and Arilla, E., β-Adrenergic regulation of the somatostatinergic system in rat hippocampus, 27
Lui, P.-W., see Fu, M.-J., 199
Lustig, H.S., see von B. Ahern, K., 211
Lutz-Bucher, B. and Koch, B., Evidence for an inhibitory effect of nitric oxides on neuropeptide secretion from isolated neural lobe of the rat pituitary gland, 48

Maione, S., Berrino, L., Pizzirusso, A., Leyva, J., Stella, L. and Rossi, F., Evidence that arcaine increases the *N*-methyl-D-aspartate-induced cardiovascular effects into the periaqueductal gray area of anesthetized rats, 164
Martínez-Serrano, A., see Blanco, P., 5
Maxwell, G.D., see Sun, Y., 1
McKenzie, J.C., see Young, J.K., 13
Meyer, W., NADPH diaphorase (nitric oxide synthase) in the central nervous system of spiders (Arachnida: Araneida), 105
Milekhina, O.N., see Supin, A.Y., 195
Misawa, M., see Kamei, J., 141
Mishina, M., see Watanabe, M., 183
Moghaddam, B., see Youngren, K.D., 41
Mogi, M., Harada, M., Riederer, P., Narabayashi, H., Fujita, K. and Nagatsu, T., Tumor necrosis factor-α (TNF-α) increases both in the brain and in the cerebrospinal fluid from parkinsonian patients, 208
Montesi, M.P., see Mortilla, M., 45
Morita, S., see Yamada, K., 191
Mortilla, M., Amaducci, L., Bruni, A., Montesi, M.P., Trubnikov, A., De Cataldo, S., Pallanti, S., Pazzagli, A., Grecu, L., Servi, P., Zazzo, M. and Sorbi, S., Absence of APP713 mutation in Italian and Russian families with schizophrenia, 45
Motohashi, O., Suzuki, M., Yanai, N., Umezawa, K., Shida, N., Shirane, R. and Yoshimoto, T., Primary culture of human leptomeningeal cells in serum-free medium, 122
Murase, K., see Sugitani, M., 215

Nabeshima, T., see Yamada, K., 191
Nagase, H., see Kamei, J., 141
Nagatsu, I., see Fujii, T., 161
Nagatsu, T., see Mogi, M., 208
Nakajima, S., see Farkas, R.H., 153
Nakanishi, H., see Ueda, Y., 203
Narabayashi, H., see Mogi, M., 208
Nishikawa, T., see Shirayama, Y., 219
Nistri, A., see Fisher, N.D., 84

O'Hara, B.F., see Bitting, L., 117
Ohm, T.G., see Derouiche, A., 179
Onoda, N., see Sugitani, M., 215

Pallanti, S., see Mortilla, M., 45
Pang, C.S., Tsang, K.F.L., Brown, G.M. and Pang, S.F., Specific 2-[^{125}I]iodomelatonin binding sites in the duck adrenal gland, 55
Pang, S.F., see Pang, C.S., 55
Park, M.J., Chung, K. and Chung, J.M., Immunohistochemical evidence for sprouting of ventral root afferents after neonatal sciatic neurectomy in the rat, 125
Parsons, C.G., see Herrero, J.F., 37
Paterno, G.D., see Reynolds, J.N., 129
Pazzagli, A., see Mortilla, M., 45
Pereira, R., see Blanco, P., 5
Pizzirusso, A., see Maione, S., 164
Popov, V.V., see Supin, A.Y., 195
Pouplard-Barthelaix, A., see Dehaut, F., 59
Prasad, A., see Reynolds, J.N., 129
Pritz, M.B. and Stritzel, M.E., Glutamic acid decarboxylase immunoreactivity in some dorsal thalamic nuclei in *Crocodilia*, 109

Reid, G., see Koh, D.-S., 167
Reymann, K.G., see Behnisch, T., 23
Reynolds, J.N., Ryan, P.J., Prasad, A. and Paterno, G.D., Neurons derived from embryonal carcinoma (P19) cells express multiple

$GABA_A$ receptor subunits and fully functional $GABA_A$ receptors, 129
Richards, H.K., see Wells, T., 93
Riederer, P., see Mogi, M., 208
Rimondini, R., see Gandolfi, O., 113
Roe, A.W., see Garraghty, P.E., 223
Rossi, F., see Maione, S., 164
Roth, R.H., see Youngren, K.D., 41
Ryan, P.J., see Reynolds, J.N., 129

Saguer, M., see Baron, R., 97
Sakai, M., see Fujii, T., 161
Satrústegui, J., see Blanco, P., 5
Schlingensiepen, K.-H., see Herms, J., 171
Schlingensiepen, R., see Herms, J., 171
Schwartz, L., see Avidor, T., 133
Servi, P., see Mortilla, M., 45
Shida, N., see Motohashi, O., 122
Shirane, R., see Motohashi, O., 122
Shirayama, Y., Nishikawa, T. and Takahashi, K., Differential effects of repeated dl-pentazocine treatment on σ binding sites in discrete brain areas of the rat, 219
Sindou, P., see Diop, A.G., 187
Smith, G.D., Wiseman, J., Harrison, S.M., Elliott, P.J. and Birch, P.J., Pre treatment with MK-801, a non-competitive NMDA antagonist, prevents development of mechanical hyperalgesia in a rat model of chronic neuropathy, but not in a model of chronic inflammation, 79
Sorbi, S., see Mortilla, M., 45
Stella, L., see Maione, S., 164
Stojilkovic, S.S., see Krsmanovic, L.Z., 33
Stritzel, M.E., see Pritz, M.B., 109
Sugai, T., see Sugitani, M., 215
Sugitani, M., Sugai, T., Tanifuji, M., Murase, K. and Onoda, N., Optical imaging of the in vitro guinea pig piriform cortex activity using a voltage-sensitive dye, 215
Sun, Y. and Maxwell, G.D., Ciliary neurotrophic factor (CNTF) has a dose-dependent biphasic effect on the number of adrenergic cells which develop in avian trunk neural crest cultures, 1
Supin, A.Y., Popov, V.V. and Milekhina, O.N., Frequency resolving power of the human's hearing, 195
Sur, M., see Garraghty, P.E., 223
Sutin, E.L., see Bitting, L., 117
Suzuki, M., see Motohashi, O., 122
Szallasi, A. and Goso, C., Characterization by [^{3}H]resiniferatoxin binding of a human vanilloid (capsaicin) receptor in post-mortem spinal cord, 101
Szücs, K., Ledesma, M.D., Dombrádi, V., Gergely, P., Avila, J. and Friedrich, P., Dephosphorylation of tau protein from Alzheimer's disease patients, 175

Takahashi, K., see Shirayama, Y., 219
Tanaka, R., see Watanabe, K., 9
Tanifuji, M., see Sugitani, M., 215
Teraoka, T., see Yamada, K., 191
Trubnikov, A., see Mortilla, M., 45
Tsai, H.-F., Kuo, T.B.J., Chan, J.Y.H. and Chan, S.H.H., Interaction between neuronal responses to nociception and hypertension in the nucleus reticularis gigantocellularis of the rat, 137
Tsang, K.F.L., see Pang, C.S., 55

Ueda, Y., Walsh, E., Nakanishi, H. and Yoshida, K., A colorimetric assay method for the evaluation of neurotrophic activity in vitro, 203
Umezawa, K., see Motohashi, O., 122

Valsecchi, B., see Gandolfi, O., 113
Verbalis, J.G., see Dohanics, J., 51
Villalba, M., see Blanco, P., 5
Virmani, M.A., see Krsmanovic, L.Z., 33
Vogel, W., see Koh, D.-S., 167
von B. Ahern, K., Lustig, H.S. and Greenberg, D.A., Enhancement of NMDA toxicity and calcium responses by chronic exposure of cultured cortical neurons to ethanol, 211

Wall, P.D., Impulses in the rostral branch of primary afferents in rat dorsal columns travel faster than those in the caudal branch, 75
Walsh, E., see Ueda, Y., 203
Watanabe, K., Fujie, S. and Tanaka, R., Effects on EMG activities of elbow joint muscles by rapid head rotation movement in horizontal plane in man, 9
Watanabe, M., Mishina, M. and Inoue, Y., Distinct gene expression of the *N*-methyl-D-aspartate receptor channel subunit in peripheral neurons of the mouse sensory ganglia and adrenal gland, 183
Watson, F.L., see Bitting, L., 117
Wells, T., Achaaban, M.R., Richards, H.K. and Forsling, M.L., Changes in neuronal metabolism in hypothalamic periventricular regions following acute alteration of cerebrospinal fluid composition in conscious rats, 93
Wiseman, J., see Smith, G.D., 79

Yamada, K., Teraoka, T., Morita, S., Hasegawa, T. and Nabeshima, T., ω-Conotoxin GVIA inhibits the methylphenidate-induced but not methamphetamine-induced behavior, 191
Yanai, N., see Motohashi, O., 122
Yáñez, J. and Anadón, R., Are the dopaminergic cells of the lamprey retina interplexiform cells? A dopamine, tyrosine hydroxylase and dopamine β-hydroxylase immuno- cytochemical study, 63
Yang, Y.L., see Lee, E.H.Y., 149
Yoshida, K., see Ueda, Y., 203
Yoshimoto, T., see Motohashi, O., 122
Young, J.K., McKenzie, J.C., Brady, L.S. and Herkenham, M., Hypothalamic lesions increase levels of neuropeptide Y mRNA in the arcuate nucleus of mice, 13
Youngren, K.D., Moghaddam, B., Bunney, B.S. and Roth, R.H., Preferential activation of dopamine overflow in prefrontal cortex produced by chronic clozapine treatment, 41

Zazzo, M., see Mortilla, M., 45
Zurmöhle, U., see Herms, J., 171